

The ABI on GOES-R

Timothy J. Schmit

NOAA/NESDIS/Satellite Applications and Research

Advanced Satellite Products Branch (ASPB)

Kaba Bah, Mathew M. Gunshor, Jun Li, Scott Bachmeier, William Straka, etc.

CIMSS, Madison, WI

James J. Gurka, Steve Goodman, etc.

GOES-R Program Office





Eighth Annual Symposium on Future Operational Environmental Satellite Systems

New Orleans, LA 24 January 2011





Also Thanks to...

- Achtor, Tom; Ackerman, Steve; Antonelli, Paolo; Aune, Bob; Baggett, Kevin; Baum, Bryan; Ellrod, Gary; Feltz, Joleen; Feltz, Wayne; Frey, Rich; Griffin, Michael K.; Gumley, Liam; Heymann, Roger; Hillger, Don; Huang, Allen; Key, Jeff; Knuteson, Bob; Mecikalski, John; Menzel, Paul; Moeller, Chris; Mosher, Fred; Nelson, James; Nasiri, Shaima; Olander, Tim; Plokhenko, Youri; Prins, Elaine; Rabin, Bob; Revercomb, Hank; Schmidt, Chris; Schreiner, Tony; Seemann-Wetzel, Suzanne; Sieglaff, Justin; Strabala, Kathy; Sun, Fengying; Tobin, Dave; Velden, Chris; Wade, Gary; Whittaker, Tom; Woolf, Hal, Jason Oktin, etc.
- Mitch Goldberg, AWG co-chairs, AWG Leads, Jaime Daniels, Walter Wolf, GPO, Jordan Gerth, Chian-Yi Liu, Jason Otkin, Thomas Greenwald, Monica Coakley, GOES-R flight/ground, Bill Smith, ASPB, PG, SSEC data center, CWG, NASA, ITT, etc.
- You!

Outline

- Current GOES Imager and Sounder
 - GOES-14/15
- ABI (Advanced Baseline Imager)
 - Temporal
 - Spatial
 - Spectral
- Summary
 - More information
 - Questions



GOES-15

GOES-13/14/15 have similar instruments to GOES-8-12, but on a different spacecraft bus.

Spring and fall eclipse outages will be avoided by larger onboard batteries.


GOES-8/12

Improved navigation


Improved radiometrics

Similar stray light to GOES-13/14


Operational on Dec 6th, 2011



Improved GOES-15 WV imagery




GOES-15 imagery




Outline

- Current GOES Imager and Sounder
 - GOES-14/15
- ABI (Advanced Baseline Imager)
 - Temporal
 - Spatial
 - Spectral
- Summary
 - More information
 - Questions



The Advanced Baseline Imager:


	ABI	Current
Spectral Coverage	16 bands	5 bands
	TO Darius	J barius
Spatial resolution		
0.64 μm Visible	0.5 km	Approx. 1 km
Other Visible/near-IR	1.0 km	n/a
Bands (>2 μm)	2 km	Approx. 4 km
Spatial coverage		
Full disk	4 per hour	Scheduled (3 hrly)
CONUS	12 per hour	~4 per hour
Mesoscale	Every 30 sec	n/a
Visible (reflective bands)		
On-orbit calibration	Yes	No 8



ABI scans about 5 times faster than the current GOES imager


There are two anticipated scan modes for the ABI:


- Full disk images every 15 minutes + 5 min CONUS images + mesoscale.₉ or - Full disk every 5 minutes.



ABI can offer Continental US images every 5 minutes for routine monitoring of a wide range of events (storms, dust, clouds, fires, winds, etc).


This is every 15 or 30 minutes with the current GOES in routine mode.





(thunderstorms, hurricanes, fires, etc). Or two regions every 60 seconds.

GOES-14: Sample "1-min" imagery





- Visible data from the GOES-14 NOAA Science Test, lead by Hillger and Schmit 13
- Can these type loops validate meso-scale models?

ABI to Imager Noise Comparison

* From ITT at GUC7

ABI				GOES			-12	-15
#	Freq. (um)	Spec	Worst Case Estimate*	#	Freq (um)	Spec	Measured (PLT)	Measured (PLT)
7	3.9	0.1	0.10	2	3.9	1.4	0.130	0.063
8	6.185	0.1	0.06					
9	6.95	0.1	0.09	3	6.x	1.0	0.15	0.17
10	7.34	0.1	0.11					
11	8.5	0.1	0.04					
12	9.61	0.1	0.04					
13	10.35	0.1	0.05					
14	11.2	0.1	0.04	4	10.7	0.35	0.11	0.06
15	12.3	0.1	0.05	5	12.0	0.35	-	-
16	13.3	0.3	0.14	6	13.3	0.32	0.19	0.13






ABI Visible/Near-IR Bands

Future GOES imager (ABI) band	Wavelength range (µm)	Central wavelength (µm)	Nominal subsatellite IGFOV (km)	Sample use
Ι	0.45-0.49	0.47	I	Daytime aerosol over land, coastal water mapping
2	0.59-0.69	0.64	0.5	Daytime clouds fog, inso- lation, winds
3	0.846-0.885	0.865	I	Daytime vegetation/burn scar and aerosol over water, winds
4	1.371-1.386	1.378	2	Daytime cirrus cloud
5	1.58–1.64	1.61	I	Daytime cloud-top phase and particle size, snow
6	2.225–2.275	2.25	2	Daytime land/cloud properties, particle size, vegetation, snow


Visible and near-IR channels on the ABI




ABI IR Bands

7	3.80-4.00	3.90	2	Surface and cloud, fog at night, fire, winds	
8	5.77–6.6	6.19	2	High-level atmospheric water vapor, winds, rainfall	
9	6.75–7.15	6.95	2	Midlevel atmospheric water vapor, winds, rainfall	
10	7.24–7.44	7.34	2	Lower-level water vapor, winds, and SO ₂	
П	8.3–8.7	8.5	2	Total water for stability, cloud phase, dust, SO ₂ rainfall	
12	9.42–9.8	9.61	2	Total ozone, turbulence, and winds	
13	10.1-10.6	10.35	2	Surface and cloud	
14	10.8–11.6	11.2	2	lmagery, SST, clouds, rainfall	
15	11.8–12.8	12.3	2	Total water, ash, and SST	
16	13.0-13.6	13.3	2	Air temperature, cloud heights and amounts	

ABI IR Weighting Functions




The IR channels on the ABI



ABI has many more bands than the current operational GOES imagers.


ABI band 1 (0.47 μm) reflectance 2008-06-26



These NWP model simulations were performed on the 'cobalt' supercomputer at the National Center for Supercomputing Applications at the University of Illinois.


Band 1: Daytime "Blue" band - aerosols, solar insolation, snow cover

ABI band 2 (0.64 µm) reflectance 2008-06-26




Band 2: Daytime "Red" band – clouds, cloud-mask, optical depth, winds, etc. ²³

ABI band 3 (0.87 μm) reflectance 2008-06-26




Band 3: Daytime "Veggie" band – NDVI, solar insolation, snow cover

ABI band 4 (1.38 μm) reflectance 2008-06-26




Band 4: Daytime "Cirrus" band - cloud mask, aerosol detection

ABI band 5 (1.61 μm) reflectance 2008-06-26




Band 5: Daytime "Snow" band – snow cover, cloud mask, etc.

ABI band 6 (2.25 μm) reflectance 2008-06-26




Band 6: Daytime "Cloud-top phase" band – cloud particle size, snow cover

ABI band 7 (3.90 μm) BT (K) 2008-06-26




Band 7: Shortwave IR window band - fog, fires, winds, SST, etc.

ABI band 8 (6.19 μm) BT (K) 2008-06-26




Band 8: Upper-level tropospheric water vapor band – moisture, flow, winds

ABI band 9 (6.95 μm) BT (K) 2008-06-26




Band 9: Upper/mid-level tropospheric water vapor band – moisture, flow, winds ³⁰

ABI band 10 (7.34 μ m) BT (K) 2008-06-26




Band 10: Lower mid-level tropospheric water vapor band— moisture, flow, winds ³¹

ABI band 11 (8.5 μm) BT (K) 2008-06-26




Band 11: "Cloud-top phase" band – SO₂, dust, SST, stability indices, etc.

ABI band 12 (9.6 μm) BT (K) 2008-06-26




Band 12: "Ozone" band

ABI band 13 (10.4 μm) BT (K) 2008-06-26




Band 13: "Clean" IR longwave window band – imagery, TPW, etc.

ABI band 14 (11.2 μm) BT (K) 2008-06-26




Band 14: IR longwave window band – many cloud parameters, SST, snow cover 35

ABI band 15 (12.3 μm) BT (K) 2008-06-26




Band 15: "Dirty" IR longwave window band – many cloud parameters, TPW ³⁶

ABI band 16 (13.3 μm) BT (K) 2008-06-26



Band 16: "CO2" longwave IR band – cloud height/pressure, stability indices ³⁷

ABI in AWIPS (via netCDF)




- 38

J. Oktin et al., CIMSS




Visualization ("decision aid")


 "True Color" with "synthetic" green from ABI simulated data (from CIMSS); image from Don Hillger, RAMMB.



Simulated ABI band – NSSL WRF



WRF Model-Derived Proxy Radiance Datasets (UW CIMSS)



Which is which – Proxy/Observed or Observed/Proxy SEVIRI 10.8 μm Tb?



GOES-R Products



Baseline Products

Advanced Baseline Imager (ABI)

Aerosol Detection (Including Smoke and Dust)

Aerosol Optical Depth (AOD)

Clear Sky Masks

Cloud and Moisture Imagery

Cloud Optical Depth

Cloud Particle Size Distribution

Cloud Top Height

Cloud Top Phase

Cloud Top Pressure

Cloud Top Temperature

Derived Motion Winds

Derived Stability Indices

Downward Shortwave Radiation: Surface

Fire/Hot Spot Characterization

Hurricane Intensity Estimation

Land Surface Temperature (Skin)

Legacy Vertical Moisture Profile

Legacy Vertical Temperature Profile

Radiances

Rainfall Rate/QPE

Reflected Shortwave Radiation: TOA

Sea Surface Temperature (Skin)

Snow Cover

Total Precipitable Water

Volcanic Ash: Detection and Height

Geostationary Lightning Mapper (GLM)

Lightning Detection: Events, Groups & Flashes

Space Environment In-Situ Suite (SEISS)

Energetic Heavy Ions

Magnetospheric Electrons & Protons: Low

Energy

Magnetospheric Electrons: Med & High Energy

Magnetospheric Protons: Med & High Energy

Solar and Galactic Protons

Magnetometer (MAG)

Geomagnetic Field

Extreme Ultraviolet and X-ray Irradiance Suite (EXIS)

Solar Flux: EUV

Solar Flux: X-ray Irradiance

Solar Ultraviolet Imager (SUVI)

Solar EUV Imagery

Future Capabilities

Advanced Baseline Imager (ABI)

Absorbed Shortwave Radiation: Surface

Aerosol Particle Size

Aircraft Icing Threat

Cloud Ice Water Path

Cloud Layers/Heights

Cloud Liquid Water

Cloud Type

Convective Initiation

Currents

Currents: Offshore

Downward Longwave Radiation: Surface Enhanced "V"/Overshooting Top Detection

Flood/Standing Water

Ice Cover

Low Cloud and Fog

Ozone Total

Probability of Rainfall

Rainfall Potential

Sea and Lake Ice: Age

Sea and Lake Ice: Concentration

Sea and Lake Ice: Motion

Snow Depth (Over Plains)

SO₂ Detection Surface Albedo

Surface Emissivity

Tropopause Folding Turbulence Prediction

Upward Longwave Radiation: Surface

Upward Longwave Radiation: TOA


Vegetation Fraction: Green

Vegetation Index

Visibility

Outline

- Current GOES Imager and Sounder
 - GOES-14/15
- ABI (Advanced Baseline Imager)
 - Temporal
 - Spatial
 - Spectral
- Summary
 - More information
 - Questions



Approximate spectral and spatial resolutions of US GOES Imagers

	~ Band Center (um)	GOES-6/7	GOES-8/11	GOES-12/13	GOES-14/15	GOES-R+
VISIDIE	0.47					
VIS	0.64	*	•			
Near-IK	0.86					
	1.6	Bo.				
	1.38	B 0.				
	2.2					
	3.9	<i>-</i>	×			×
	6.2					
	6.5/6.7/7	14km		4		2
	7.3	"MSI mode"				
IIIITareu	8.5	······				×
TUID	9.7					<u>×</u>
	10.35					<u>×</u>
	11.2	,	×	×		<u>×</u>
	12.3					
	13.3				4	×

Summary



- The ABI on GOES-R will improve over the current instruments (spectrally, spatially and temporally), plus improved image navigation and registration and radiometer performance.
- These improvements will greatly assist a host of applications and new products.
- Contact information:
 - tim.j.schmit@noaa.gov

More information

GOES-R:

- http://www.goes-r.gov
- http://www.meted.ucar.edu/index.htm
- http://cimss.ssec.wisc.edu/goes_r/proving-ground.html

GOES and NASA:

- http://goespoes.gsfc.nasa.gov/goes/index.html
- http://goes.gsfc.nasa.gov/text/goes.databookn.html

UW/SSEC/CIMSS/ASPB:

- http://cimss.ssec.wisc.edu/goes_r/provingground/nssl_abi/nssl_abi_rt.html
- http://cimss.ssec.wisc.edu/goes_r/awg/proxy/nwp/
- http://cimss.ssec.wisc.edu/goes/abi/
- http://cimss.ssec.wisc.edu/goes/abi/wf
- http://cimss.ssec.wisc.edu/goes/blog/
- http://www.ssec.wisc.edu/data/geo/





Acknowledgements



- The authors would like to thank the entire GOES-R team (especially the GOES-R Program Office); within the government, industry and academia.
- The views, opinions, and findings contained in this presentation are those of the author and should not be construed as an official National Oceanic and Atmospheric Administration or U.S. Government position, policy, or decision.





